



International Baccalaureate
Baccalauréat International
Bachillerato Internacional

Edukraft[®]
Crafting Your Child's Future
IB ALEVEL IGCSE SAT ACT

Diploma Programme

Mathematics HL and further mathematics HL formula booklet

For use during the course and in the examinations

First examinations 2014

Edukraft®

Crafting Your Child's Future

IB

A LEVEL

IGCSE

SAT

ACT

Contents

Prior learning	2
Core	3
Topic 1: Algebra	3
Topic 2: Functions and equations	4
Topic 3: Circular functions and trigonometry	4
Topic 4: Vectors	5
Topic 5: Statistics and probability	6
Topic 6: Calculus	8
Options	10
Topic 7: Statistics and probability	10
Further mathematics HL topic 3	
Topic 8: Sets, relations and groups	11
Further mathematics HL topic 4	
Topic 9: Calculus	11
Further mathematics HL topic 5	
Topic 10: Discrete mathematics	12
Further mathematics HL topic 6	
Formulae for distributions	13
Topics 5.6, 5.7, 7.1, further mathematics HL topic 3.1	
Discrete distributions	13
Continuous distributions	13
Further mathematics	14
Topic 1: Linear algebra	14

Prior learning

Area of a parallelogram	$A = b \times h$, where b is the base, h is the height
Area of a triangle	$A = \frac{1}{2}(b \times h)$, where b is the base, h is the height
Area of a trapezium	$A = \frac{1}{2}(a + b)h$, where a and b are the parallel sides, h is the height
Area of a circle	$A = \pi r^2$, where r is the radius
Circumference of a circle	$C = 2\pi r$, where r is the radius
Volume of a pyramid	$V = \frac{1}{3}(\text{area of base} \times \text{vertical height})$
Volume of a cuboid	$V = l \times w \times h$, where l is the length, w is the width, h is the height
Volume of a cylinder	$V = \pi r^2 h$, where r is the radius, h is the height
Area of the curved surface of a cylinder	$A = 2\pi r h$, where r is the radius, h is the height
Volume of a sphere	$V = \frac{4}{3}\pi r^3$, where r is the radius
Volume of a cone	$V = \frac{1}{3}\pi r^2 h$, where r is the radius, h is the height
Distance between two points (x_1, y_1) and (x_2, y_2)	$d = \sqrt{(x_1 - x_2)^2 + (y_1 - y_2)^2}$
Coordinates of the midpoint of a line segment with endpoints (x_1, y_1) and (x_2, y_2)	$\left(\frac{x_1 + x_2}{2}, \frac{y_1 + y_2}{2}\right)$
Solutions of a quadratic equation	The solutions of $ax^2 + bx + c = 0$ are $x = \frac{-b \pm \sqrt{b^2 - 4ac}}{2a}$

Topic I: Algebra

1.1	<p>The n^{th} term of an arithmetic sequence</p> <p>The sum of n terms of an arithmetic sequence</p> <p>The n^{th} term of a geometric sequence</p> <p>The sum of n terms of a finite geometric sequence</p> <p>The sum of an infinite geometric sequence</p>	$u_n = u_1 + (n-1)d$ $S_n = \frac{n}{2}(2u_1 + (n-1)d) = \frac{n}{2}(u_1 + u_n)$ $u_n = u_1 r^{n-1}$ $S_n = \frac{u_1(r^n - 1)}{r - 1} = \frac{u_1(1 - r^n)}{1 - r}, r \neq 1$ $S_\infty = \frac{u_1}{1 - r}, r < 1$
1.2	Exponents and logarithms	$a^x = b \Leftrightarrow x = \log_a b, \text{ where } a > 0, b > 0, a \neq 1$ $a^x = e^{x \ln a}$ $\log_a a^x = x = a^{\log_a x}$ $\log_b a = \frac{\log_c a}{\log_c b}$
1.3	<p>Combinations</p> <p>Permutations</p> <p>Binomial theorem</p>	$\binom{n}{r} = \frac{n!}{r!(n-r)!}$ ${}_n P_r = \frac{n!}{(n-r)!}$ $(a+b)^n = a^n + \binom{n}{1} a^{n-1} b + \dots + \binom{n}{r} a^{n-r} b^r + \dots + b^n$
1.5	Complex numbers	$z = a + ib = r(\cos \theta + i \sin \theta) = re^{i\theta} = r \operatorname{cis} \theta$
1.7	De Moivre's theorem	$[r(\cos \theta + i \sin \theta)]^n = r^n (\cos n\theta + i \sin n\theta) = r^n e^{in\theta} = r^n \operatorname{cis} n\theta$

Topic 2: Functions and equations

2.5	Axis of symmetry of the graph of a quadratic function	$f(x) = ax^2 + bx + c \Rightarrow$ axis of symmetry $x = -\frac{b}{2a}$
2.6	Discriminant	$\Delta = b^2 - 4ac$

Topic 3: Circular functions and trigonometry

3.1	Length of an arc Area of a sector	$l = \theta r$, where θ is the angle measured in radians, r is the radius $A = \frac{1}{2}\theta r^2$, where θ is the angle measured in radians, r is the radius
3.2	Identities Pythagorean identities	$\tan \theta = \frac{\sin \theta}{\cos \theta}$ $\sec \theta = \frac{1}{\cos \theta}$ $\operatorname{cosec} \theta = \frac{1}{\sin \theta}$ $\cos^2 \theta + \sin^2 \theta = 1$ $1 + \tan^2 \theta = \sec^2 \theta$ $1 + \cot^2 \theta = \operatorname{csc}^2 \theta$
3.3	Compound angle identities Double angle identities	$\sin(A \pm B) = \sin A \cos B \pm \cos A \sin B$ $\cos(A \pm B) = \cos A \cos B \mp \sin A \sin B$ $\tan(A \pm B) = \frac{\tan A \pm \tan B}{1 \mp \tan A \tan B}$ $\sin 2\theta = 2 \sin \theta \cos \theta$ $\cos 2\theta = \cos^2 \theta - \sin^2 \theta = 2 \cos^2 \theta - 1 = 1 - 2 \sin^2 \theta$ $\tan 2\theta = \frac{2 \tan \theta}{1 - \tan^2 \theta}$

3.7	Cosine rule	$c^2 = a^2 + b^2 - 2ab \cos C$; $\cos C = \frac{a^2 + b^2 - c^2}{2ab}$
	Sine rule	$\frac{a}{\sin A} = \frac{b}{\sin B} = \frac{c}{\sin C}$
	Area of a triangle	$A = \frac{1}{2} ab \sin C$

Topic 4: Vectors

4.1	Magnitude of a vector	$ \mathbf{v} = \sqrt{v_1^2 + v_2^2 + v_3^2}$, where $\mathbf{v} = \begin{pmatrix} v_1 \\ v_2 \\ v_3 \end{pmatrix}$
	Distance between two points (x_1, y_1, z_1) and (x_2, y_2, z_2)	$d = \sqrt{(x_1 - x_2)^2 + (y_1 - y_2)^2 + (z_1 - z_2)^2}$
	Coordinates of the midpoint of a line segment with endpoints (x_1, y_1, z_1) , (x_2, y_2, z_2)	$\left(\frac{x_1 + x_2}{2}, \frac{y_1 + y_2}{2}, \frac{z_1 + z_2}{2} \right)$
4.2	Scalar product	$\mathbf{v} \cdot \mathbf{w} = \mathbf{v} \mathbf{w} \cos \theta$, where θ is the angle between \mathbf{v} and \mathbf{w}
	Angle between two vectors	$\mathbf{v} \cdot \mathbf{w} = v_1 w_1 + v_2 w_2 + v_3 w_3$, where $\mathbf{v} = \begin{pmatrix} v_1 \\ v_2 \\ v_3 \end{pmatrix}$, $\mathbf{w} = \begin{pmatrix} w_1 \\ w_2 \\ w_3 \end{pmatrix}$ $\cos \theta = \frac{v_1 w_1 + v_2 w_2 + v_3 w_3}{ \mathbf{v} \mathbf{w} }$
4.3	Vector equation of a line	$\mathbf{r} = \mathbf{a} + \lambda \mathbf{b}$
	Parametric form of the equation of a line	$x = x_0 + \lambda l$, $y = y_0 + \lambda m$, $z = z_0 + \lambda n$
	Cartesian equations of a line	$\frac{x - x_0}{l} = \frac{y - y_0}{m} = \frac{z - z_0}{n}$

4.5	Vector product Area of a triangle	$\mathbf{v} \times \mathbf{w} = \begin{pmatrix} v_2 w_3 - v_3 w_2 \\ v_3 w_1 - v_1 w_3 \\ v_1 w_2 - v_2 w_1 \end{pmatrix} \text{ where } \mathbf{v} = \begin{pmatrix} v_1 \\ v_2 \\ v_3 \end{pmatrix}, \mathbf{w} = \begin{pmatrix} w_1 \\ w_2 \\ w_3 \end{pmatrix}$ $ \mathbf{v} \times \mathbf{w} = \mathbf{v} \mathbf{w} \sin \theta, \text{ where } \theta \text{ is the angle between } \mathbf{v} \text{ and } \mathbf{w}$ $A = \frac{1}{2} \mathbf{v} \times \mathbf{w} \text{ where } \mathbf{v} \text{ and } \mathbf{w} \text{ form two sides of a triangle}$
4.6	Vector equation of a plane Equation of a plane (using the normal vector) Cartesian equation of a plane	$\mathbf{r} = \mathbf{a} + \lambda \mathbf{b} + \mu \mathbf{c}$ $\mathbf{r} \cdot \mathbf{n} = \mathbf{a} \cdot \mathbf{n}$ $ax + by + cz = d$

Topic 5: Statistics and probability

5.1	Population parameters Mean μ Variance σ^2 Standard deviation σ	$\text{Let } n = \sum_{i=1}^k f_i$ $\mu = \frac{\sum_{i=1}^k f_i x_i}{n}$ $\sigma^2 = \frac{\sum_{i=1}^k f_i (x_i - \mu)^2}{n} = \frac{\sum_{i=1}^k f_i x_i^2}{n} - \mu^2$ $\sigma = \sqrt{\frac{\sum_{i=1}^k f_i (x_i - \mu)^2}{n}}$
5.2	Probability of an event A Complementary events	$P(A) = \frac{n(A)}{n(U)}$ $P(A) + P(A') = 1$
5.3	Combined events Mutually exclusive events	$P(A \cup B) = P(A) + P(B) - P(A \cap B)$ $P(A \cup B) = P(A) + P(B)$

<p>5.4</p>	<p>Conditional probability</p> <p>Independent events</p> <p>Bayes' theorem</p>	$P(A B) = \frac{P(A \cap B)}{P(B)}$ $P(A \cap B) = P(A)P(B)$ $P(B A) = \frac{P(B)P(A B)}{P(B)P(A B) + P(B')P(A B')}$ $P(B_i A) = \frac{P(B_i)P(A B_i)}{P(B_1)P(A B_1) + P(B_2)P(A B_2) + P(B_3)P(A B_3)}$
<p>5.5</p>	<p>Expected value of a discrete random variable X</p> <p>Expected value of a continuous random variable X</p> <p>Variance</p> <p>Variance of a discrete random variable X</p> <p>Variance of a continuous random variable X</p>	$E(X) = \mu = \sum x P(X = x)$ $E(X) = \mu = \int_{-\infty}^{\infty} x f(x) dx$ $\text{Var}(X) = E(X - \mu)^2 = E(X^2) - [E(X)]^2$ $\text{Var}(X) = \sum (x - \mu)^2 P(X = x) = \sum x^2 P(X = x) - \mu^2$ $\text{Var}(X) = \int_{-\infty}^{\infty} (x - \mu)^2 f(x) dx = \int_{-\infty}^{\infty} x^2 f(x) dx - \mu^2$
<p>5.6</p>	<p>Binomial distribution</p> <p>Mean</p> <p>Variance</p> <p>Poisson distribution</p> <p>Mean</p> <p>Variance</p>	$X \sim B(n, p) \Rightarrow P(X = x) = \binom{n}{x} p^x (1-p)^{n-x}, \quad x = 0, 1, \dots, n$ $E(X) = np$ $\text{Var}(X) = np(1-p)$ $X \sim \text{Po}(m) \Rightarrow P(X = x) = \frac{m^x e^{-m}}{x!}, \quad x = 0, 1, 2, \dots$ $E(X) = m$ $\text{Var}(X) = m$
<p>5.7</p>	<p>Standardized normal variable</p>	$z = \frac{x - \mu}{\sigma}$

Topic 6: Calculus

6.1	Derivative of $f(x)$	$y = f(x) \Rightarrow \frac{dy}{dx} = f'(x) = \lim_{h \rightarrow 0} \left(\frac{f(x+h) - f(x)}{h} \right)$
6.2	Derivative of x^n	$f(x) = x^n \Rightarrow f'(x) = nx^{n-1}$
	Derivative of $\sin x$	$f(x) = \sin x \Rightarrow f'(x) = \cos x$
	Derivative of $\cos x$	$f(x) = \cos x \Rightarrow f'(x) = -\sin x$
	Derivative of $\tan x$	$f(x) = \tan x \Rightarrow f'(x) = \sec^2 x$
	Derivative of e^x	$f(x) = e^x \Rightarrow f'(x) = e^x$
	Derivative of $\ln x$	$f(x) = \ln x \Rightarrow f'(x) = \frac{1}{x}$
	Derivative of $\sec x$	$f(x) = \sec x \Rightarrow f'(x) = \sec x \tan x$
	Derivative of $\csc x$	$f(x) = \csc x \Rightarrow f'(x) = -\csc x \cot x$
	Derivative of $\cot x$	$f(x) = \cot x \Rightarrow f'(x) = -\csc^2 x$
	Derivative of a^x	$f(x) = a^x \Rightarrow f'(x) = a^x (\ln a)$
	Derivative of $\log_a x$	$f(x) = \log_a x \Rightarrow f'(x) = \frac{1}{x \ln a}$
	Derivative of $\arcsin x$	$f(x) = \arcsin x \Rightarrow f'(x) = \frac{1}{\sqrt{1-x^2}}$
	Derivative of $\arccos x$	$f(x) = \arccos x \Rightarrow f'(x) = -\frac{1}{\sqrt{1-x^2}}$
	Derivative of $\arctan x$	$f(x) = \arctan x \Rightarrow f'(x) = \frac{1}{1+x^2}$
	Chain rule	$y = g(u), \text{ where } u = f(x) \Rightarrow \frac{dy}{dx} = \frac{dy}{du} \times \frac{du}{dx}$
	Product rule	$y = uv \Rightarrow \frac{dy}{dx} = u \frac{dv}{dx} + v \frac{du}{dx}$
	Quotient rule	$y = \frac{u}{v} \Rightarrow \frac{dy}{dx} = \frac{v \frac{du}{dx} - u \frac{dv}{dx}}{v^2}$

6.4	Standard integrals	$\int x^n dx = \frac{x^{n+1}}{n+1} + C, \quad n \neq -1$ $\int \frac{1}{x} dx = \ln x + C$ $\int \sin x dx = -\cos x + C$ $\int \cos x dx = \sin x + C$ $\int e^x dx = e^x + C$ $\int a^x dx = \frac{1}{\ln a} a^x + C$ $\int \frac{1}{a^2 + x^2} dx = \frac{1}{a} \arctan\left(\frac{x}{a}\right) + C$ $\int \frac{1}{\sqrt{a^2 - x^2}} dx = \arcsin\left(\frac{x}{a}\right) + C, \quad x < a$
6.5	Area under a curve Volume of revolution (rotation)	$A = \int_a^b y dx \text{ or } A = \int_a^b x dy$ $V = \int_a^b \pi y^2 dx \text{ or } V = \int_a^b \pi x^2 dy$
6.7	Integration by parts	$\int u \frac{dv}{dx} dx = uv - \int v \frac{du}{dx} dx \text{ or } \int u dv = uv - \int v du$

Topic 7: Statistics and probability

Further mathematics HL topic 3

7.1 (3.1)	Probability generating function for a discrete random variable X	$G(t) = E(t^x) = \sum_x P(X = x)t^x$ $E(X) = G'(1)$ $\text{Var}(X) = G''(1) + G'(1) - (G'(1))^2$
7.2 (3.2)	Linear combinations of two independent random variables X_1, X_2	$E(a_1X_1 \pm a_2X_2) = a_1E(X_1) \pm a_2E(X_2)$ $\text{Var}(a_1X_1 \pm a_2X_2) = a_1^2 \text{Var}(X_1) + a_2^2 \text{Var}(X_2)$
7.3 (3.3)	Sample statistics Mean \bar{x} Variance s_n^2 Standard deviation s_n Unbiased estimate of population variance s_{n-1}^2	$\bar{x} = \frac{\sum_{i=1}^k f_i x_i}{n}$ $s_n^2 = \frac{\sum_{i=1}^k f_i (x_i - \bar{x})^2}{n} = \frac{\sum_{i=1}^k f_i x_i^2}{n} - \bar{x}^2$ $s_n = \sqrt{\frac{\sum_{i=1}^k f_i (x_i - \bar{x})^2}{n}}$ $s_{n-1}^2 = \frac{n}{n-1} s_n^2 = \frac{\sum_{i=1}^k f_i (x_i - \bar{x})^2}{n-1} = \frac{\sum_{i=1}^k f_i x_i^2}{n-1} - \frac{n}{n-1} \bar{x}^2$
7.5 (3.5)	Confidence intervals Mean, with known variance Mean, with unknown variance	$\bar{x} \pm z \times \frac{\sigma}{\sqrt{n}}$ $\bar{x} \pm t \times \frac{s_{n-1}}{\sqrt{n}}$
7.6 (3.6)	Test statistics Mean, with known variance	$z = \frac{\bar{x} - \mu}{\sigma / \sqrt{n}}$

	Mean, with unknown variance	$t = \frac{\bar{x} - \mu}{s_{n-1} / \sqrt{n}}$
7.7 (3.7)	Sample product moment correlation coefficient	$r = \frac{\sum_{i=1}^n x_i y_i - n\bar{x}\bar{y}}{\sqrt{\left(\sum_{i=1}^n x_i^2 - n\bar{x}^2\right)\left(\sum_{i=1}^n y_i^2 - n\bar{y}^2\right)}}$
	Test statistic for H_0 : $\rho = 0$	$t = r\sqrt{\frac{n-2}{1-r^2}}$
	Equation of regression line of x on y	$x - \bar{x} = \left(\frac{\sum_{i=1}^n x_i y_i - n\bar{x}\bar{y}}{\sum_{i=1}^n y_i^2 - n\bar{y}^2} \right) (y - \bar{y})$
	Equation of regression line of y on x	$y - \bar{y} = \left(\frac{\sum_{i=1}^n x_i y_i - n\bar{x}\bar{y}}{\sum_{i=1}^n x_i^2 - n\bar{x}^2} \right) (x - \bar{x})$

Topic 8: Sets, relations and groups

Further mathematics HL topic 4

8.1 (4.1)	De Morgan's laws	$(A \cup B)' = A' \cap B'$ $(A \cap B)' = A' \cup B'$
----------------------------	------------------	--

Topic 9: Calculus

Further mathematics HL topic 5

9.5 (5.5)	Euler's method	$y_{n+1} = y_n + h \times f(x_n, y_n); \quad x_{n+1} = x_n + h, \text{ where } h \text{ is a constant (step length)}$
	Integrating factor for $y' + P(x)y = Q(x)$	$e^{\int P(x) dx}$

9.6 (5.6)	Maclaurin series	$f(x) = f(0) + x f'(0) + \frac{x^2}{2!} f''(0) + \dots$
	Taylor series	$f(x) = f(a) + (x-a)f'(a) + \frac{(x-a)^2}{2!} f''(a) + \dots$
	Taylor approximations (with error term $R_n(x)$)	$f(x) = f(a) + (x-a)f'(a) + \dots + \frac{(x-a)^n}{n!} f^{(n)}(a) + R_n(x)$
	Lagrange form	$R_n(x) = \frac{f^{(n+1)}(c)}{(n+1)!} (x-a)^{n+1}$, where c lies between a and x
	Maclaurin series for special functions	$e^x = 1 + x + \frac{x^2}{2!} + \dots$ $\ln(1+x) = x - \frac{x^2}{2} + \frac{x^3}{3} - \dots$ $\sin x = x - \frac{x^3}{3!} + \frac{x^5}{5!} - \dots$ $\cos x = 1 - \frac{x^2}{2!} + \frac{x^4}{4!} - \dots$ $\arctan x = x - \frac{x^3}{3} + \frac{x^5}{5} - \dots$

Topic 10: Discrete mathematics

Further mathematics HL topic 6

10.7 (6.7)	Euler's formula for connected planar graphs	$v - e + f = 2$, where v is the number of vertices, e is the number of edges, f is the number of faces
	Planar, simple, connected graphs	$e \leq 3v - 6$ for $v \geq 3$ $e \leq 2v - 4$ if the graph has no triangles

Edukraft[®]
Crafting Your Child's Future

IB ALEVEL IGCSE SAT ACT

Formulae for distributions

Topics 5.6, 5.7, 7.1, further mathematics HL topic 3.1

Discrete distributions

Distribution	Notation	Probability mass function	Mean	Variance
Geometric	$X \sim \text{Geo}(p)$	pq^{x-1} for $x = 1, 2, \dots$	$\frac{1}{p}$	$\frac{q}{p^2}$
Negative binomial	$X \sim \text{NB}(r, p)$	$\binom{x-1}{r-1} p^r q^{x-r}$ for $x = r, r+1, \dots$	$\frac{r}{p}$	$\frac{rq}{p^2}$

Continuous distributions

Distribution	Notation	Probability density function	Mean	Variance
Normal	$X \sim N(\mu, \sigma^2)$	$\frac{1}{\sigma\sqrt{2\pi}} e^{-\frac{1}{2}\left(\frac{x-\mu}{\sigma}\right)^2}$	μ	σ^2

Topic I: Linear algebra

1.2	Determinant of a 2×2 matrix	$\mathbf{A} = \begin{pmatrix} a & b \\ c & d \end{pmatrix} \Rightarrow \det \mathbf{A} = \mathbf{A} = ad - bc$
	Inverse of a 2×2 matrix	$\mathbf{A} = \begin{pmatrix} a & b \\ c & d \end{pmatrix} \Rightarrow \mathbf{A}^{-1} = \frac{1}{\det \mathbf{A}} \begin{pmatrix} d & -b \\ -c & a \end{pmatrix}, ad \neq bc$
	Determinant of a 3×3 matrix	$\mathbf{A} = \begin{pmatrix} a & b & c \\ d & e & f \\ g & h & k \end{pmatrix} \Rightarrow \det \mathbf{A} = a \begin{vmatrix} e & f \\ h & k \end{vmatrix} - b \begin{vmatrix} d & f \\ g & k \end{vmatrix} + c \begin{vmatrix} d & e \\ g & h \end{vmatrix}$